






# IFRMP - Bull Trout Range

## Legend:

-  Klamath Subbasin Groups
-  Bull Trout - current distribution<sup>2</sup>
-  Bull Trout - additional historical distribution<sup>2</sup>
-  Bull Trout - Designated Critical Habitat<sup>1</sup>
-  Keno diversion area

## Data Sources:

1 - ODFW Oregon Fish Habitat Distribution Data (ODFW Data Clearinghouse):  
<https://nrimp.dfw.state.or.us/DataClearinghouse/default.aspx?p=202&XMLname=1167.xml>


2 - USFWS Critical Habitat Designations:  
<http://criticalhabitat.fws.gov/crithab/>

## About This Map:

This map was produced as part of the Klamath Integrated Fisheries Restoration and Monitoring Plan (IFRMP) and is being used as the basis for determining range overlap of focal fish species with potential project locations during the restoration prioritization phase of this project (which is only one of several prioritization criteria). Find out more about this project at <http://kbifrm.psmfc.org/>.





Note that the Keno Diversion area shown on this map indicates reassignment of these sub-watersheds from the Lost Sub-basin to the Upper Klamath Lake subbasin for the purpose of participant group discussions in this project only. This decision was made to reflect the hydrologic connectivity of the mainstem in these sub-watersheds to Upper Klamath Lake, which drains directly into this portion of the mainstem and has a significant influence on it. This 'reorganization' will ensure that discussion of restoration priorities in this segment of the mainstem occurs in the same group discussing restoration priorities for Upper Klamath Lake.

Photo: (3) USFWS licenced under CC by 2.0



# IFRMP - Fall Chinook Range


## Legend:

-  Klamath Subbasin Groups
-  Fall Chinook - current distribution <sup>2</sup>
-  Fall Chinook - additional historical distribution <sup>1,2</sup>
-  Keno Diversion area

## Data Sources:

1 - ODFW Oregon Fish Habitat Distribution Data (ODFW Data Clearinghouse):  
<https://nrimp.dfw.state.or.us/DataClearinghouse/default.aspx?p=202&XMLname=1167.xml>

2 - UC Davis - PISCES Fish Range and Occurrence Database:  
[https://pisc.es.ucdavis.edu/fish?term\\_node\\_tid\\_depth=All&term\\_node\\_tid\\_depth\\_1=All&field\\_species\\_commonname\\_value=](https://pisc.es.ucdavis.edu/fish?term_node_tid_depth=All&term_node_tid_depth_1=All&field_species_commonname_value=)







## About This Map:

This map was produced as part of the Klamath Integrated Fisheries Restoration and Monitoring Plan (IFRMP) and is being used as the basis for determining range overlap of focal fish species with potential project locations during the restoration prioritization phase of this project (which is only one of several prioritization criteria). Find out more about this project at <http://kbifrm.psmfc.org/>.

Note that the Keno Diversion area shown on this map indicates reassignment of these sub-watersheds from the Lost Sub-basin to the Upper Klamath Lake subbasin for the purpose of participant group discussions in this project only. This decision was made to reflect the hydrologic connectivity of the mainstem in these sub-watersheds to Upper Klamath Lake, which drains directly into this portion of the mainstem and has a significant influence on it. This 'reorganization' will ensure that discussion of restoration priorities in this segment of the mainstem occurs in the same group discussing restoration priorities for Upper Klamath Lake.

# IFRMP - Spring Chinook Range


## Legend:

-  Klamath Subbasin Groups
-  Spring Chinook - current distribution <sup>1</sup>
-  Spring Chinook - additional historical distribution <sup>1, 2</sup>
-  Keno diversion area

## Data Sources:

1 - UC Davis - PISCES Fish Range and Occurrence Database: [https://pisc.es.ucdavis.edu/fish?term\\_node\\_tid\\_depth=All&term\\_node\\_tid\\_depth\\_1=All&field\\_species\\_commonname\\_value=](https://pisc.es.ucdavis.edu/fish?term_node_tid_depth=All&term_node_tid_depth_1=All&field_species_commonname_value=)

2 - ODFW Oregon Fish Habitat Distribution Data (ODFW Data Clearinghouse): <https://nrimp.dfw.state.or.us/DataClearinghouse/default.aspx?p=202&XMLname=1167.xml>







## About This Map:

This map was produced as part of the Klamath Integrated Fisheries Restoration and Monitoring Plan (IFRMP) and is being used as the basis for determining range overlap of focal fish species with potential project locations during the restoration prioritization phase of this project (which is only one of several prioritization criteria). Find out more about this project at <http://kbifrm.psmfc.org/>.

Note that the Keno Diversion area shown on this map indicates reassignment of these sub-watersheds from the Lost Sub-basin to the Upper Klamath Lake subbasin for the purpose of participant group discussions in this project only. This decision was made to reflect the hydrologic connectivity of the mainstem in these sub-watersheds to Upper Klamath Lake, which drains directly into this portion of the mainstem and has a significant influence on it. This 'reorganization' will ensure that discussion of restoration priorities in this segment of the mainstem occurs in the same group discussing restoration priorities for Upper Klamath Lake.

# IFRMP - Coho Range


## Legend:

-  Klamath Subbasin Groups
-  Coho - current distribution<sup>2</sup>
-  Coho - additional historical distribution<sup>1,2</sup>
-  Keno diversion area

## Data Sources:

1 - ODFW Oregon Fish Habitat Distribution Data (ODFW Data Clearinghouse):  
<https://nrimp.dfw.state.or.us/DataClearinghouse/default.aspx?p=202&XMLname=1167.xml>

2 - UC Davis - PISCES Fish Range and Occurrence Database:  
[https://pisc.es.ucdavis.edu/fish?term\\_node\\_tid\\_depth=All&term\\_node\\_tid\\_depth\\_1=All&field\\_species\\_commonname\\_value=](https://pisc.es.ucdavis.edu/fish?term_node_tid_depth=All&term_node_tid_depth_1=All&field_species_commonname_value=)






## About This Map:

This map was produced as part of the Klamath Integrated Fisheries Restoration and Monitoring Plan (IFRMP) and is being used as the basis for determining range overlap of focal fish species with potential project locations during the restoration prioritization phase of this project (which is only one of several prioritization criteria). Find out more about this project at <http://kbifrm.psmfc.org/>.

Note that the Keno Diversion area shown on this map indicates reassignment of these sub-watersheds from the Lost Sub-basin to the Upper Klamath Lake subbasin for the purpose of participant group discussions in this project only. This decision was made to reflect the hydrologic connectivity of the mainstem in these sub-watersheds to Upper Klamath Lake, which drains directly into this portion of the mainstem and has a significant influence on it. This 'reorganization' will ensure that discussion of restoration priorities in this segment of the mainstem occurs in the same group discussing restoration priorities for Upper Klamath Lake.


# IFRMP - Eulachon Range

## Legend:

-  Klamath Subbasin Groups
-  Eulachon - current & historical distribution<sup>1</sup>
-  Keno diversion area

## Data Sources:

1 - UC Davis - PISCES Fish Range and Occurrence Database:  
[https://pisc.es.ucdavis.edu/fish?term\\_node\\_tid\\_depth=All&term\\_node\\_tid\\_depth\\_1=All&field\\_species\\_commonname\\_value=](https://pisc.es.ucdavis.edu/fish?term_node_tid_depth=All&term_node_tid_depth_1=All&field_species_commonname_value=)



## About This Map:




This map was produced as part of the Klamath Integrated Fisheries Restoration and Monitoring Plan (IFRMP) and is being used as the basis for determining range overlap of focal fish species with potential project locations during the restoration prioritization phase of this project (which is only one of several prioritization criteria). Find out more about this project at <http://kbifrm.psmfc.org/>.

Note that the Keno Diversion area shown on this map indicates reassignment of these sub-watersheds from the Lost Sub-basin to the Upper Klamath Lake subbasin for the purpose of participant group discussions in this project only. This decision was made to reflect the hydrologic connectivity of the mainstem in these sub-watersheds to Upper Klamath Lake, which drains directly into this portion of the mainstem and has a significant influence on it. This 'reorganization' will ensure that discussion of restoration priorities in this segment of the mainstem occurs in the same group discussing restoration priorities for Upper Klamath Lake.

# IFRMP - Green Sturgeon Range




## Legend:

-  Klamath Subbasin Groups
-  Green Sturgeon - current & historical distribution<sup>1</sup>
-  Keno diversion area

## Data Sources:

1 - UC Davis - PISCES Fish Range and Occurrence Database:  
[https://pisc.es.ucdavis.edu/fish?term\\_node\\_tid\\_depth=All&term\\_node\\_tid\\_depth\\_1=All&field\\_species\\_commonname\\_value=](https://pisc.es.ucdavis.edu/fish?term_node_tid_depth=All&term_node_tid_depth_1=All&field_species_commonname_value=)










## About This Map:


This map was produced as part of the Klamath Integrated Fisheries Restoration and Monitoring Plan (IFRMP) and is being used as the basis for determining range overlap of focal fish species with potential project locations during the restoration prioritization phase of this project (which is only one of several prioritization criteria). Find out more about this project at <http://kbifrm.psmfc.org/>.

Note that the Keno Diversion area shown on this map indicates reassignment of these sub-watersheds from the Lost Sub-basin to the Upper Klamath Lake subbasin for the purpose of participant group discussions in this project only. This decision was made to reflect the hydrologic connectivity of the mainstem in these sub-watersheds to Upper Klamath Lake, which drains directly into this portion of the mainstem and has a significant influence on it. This 'reorganization' will ensure that discussion of restoration priorities in this segment of the mainstem occurs in the same group discussing restoration priorities for Upper Klamath Lake.

# IFRMP - Lost River Sucker Range

## Legend:

-  Klamath Subbasin Groups
-  Keno diversion area
-  Lost River Sucker - current distribution <sup>1</sup>
-  Lost River Sucker - additional historical distribution <sup>2</sup>
-  Lost River Sucker - Designated Critical Habitat <sup>3</sup>
-  Lost River Sucker - Designated Streams
-  Lost River Sucker - Designated Lakes



## Data Sources:

- 1 - USFWS 2019 Species Status Assessment for the Endangered Lost River and Shortnose Sucker  
[https://www.researchgate.net/publication/333787292\\_Species\\_Status\\_Assessment\\_for\\_the\\_Endangered\\_Lost\\_River\\_Sucker\\_and\\_Shortnose\\_Sucker](https://www.researchgate.net/publication/333787292_Species_Status_Assessment_for_the_Endangered_Lost_River_Sucker_and_Shortnose_Sucker)
- 2 - Josh Rasmussen, USFWS, Personal communication.
- 3 - USFWS Critical Habitat Designations:  
<http://criticalhabitat.fws.gov/crithab>

## About This Map:





This map was produced as part of the Klamath Integrated Fisheries Restoration and Monitoring Plan (IFRMP) and is being used as the basis for determining range overlap of focal fish species with potential project locations during the restoration prioritization phase of this project (which is only one of several prioritization criteria). Find out more about this project at <http://kbifrm.psmfc.org/>.

Note that the Keno Diversion area shown on this map indicates reassignment of these sub-watersheds from the Lost Sub-basin to the Upper Klamath Lake subbasin for the purpose of participant group discussions in this project only. This decision was made to reflect the hydrologic connectivity of the mainstem in these sub-watersheds to Upper Klamath Lake, which drains directly into this portion of the mainstem and has a significant influence on it. This 'reorganization' will ensure that discussion of restoration priorities in this segment of the mainstem occurs in the same group discussing restoration priorities for Upper Klamath Lake.

# IFRMP - Pacific Lamprey Range




## Legend:

-  Klamath Subbasin Groups
-  Pacific Lamprey - current distribution<sup>1</sup>
-  Pacific Lamprey - additional historical distribution<sup>1</sup>
-  Keno diversion area

## Data Sources:

1 - UC Davis - PISCES Fish Range and Occurrence Database (California, with CDFW Data);  
[https://pisc.es.ucdavis.edu/fish?term\\_node\\_tid\\_depth=All&term\\_node\\_tid\\_depth\\_1=All&field\\_species\\_commonname\\_value=](https://pisc.es.ucdavis.edu/fish?term_node_tid_depth=All&term_node_tid_depth_1=All&field_species_commonname_value=)



## About This Map:





This map was produced as part of the Klamath Integrated Fisheries Restoration and Monitoring Plan (IFRMP) and is being used as the basis for determining range overlap of focal fish species with potential project locations during the restoration prioritization phase of this project (which is only one of several prioritization criteria). Find out more about this project at <http://kbifrm.psmfc.org/>.

Note that the Keno Diversion area shown on this map indicates reassignment of these sub-watersheds from the Lost Sub-basin to the Upper Klamath Lake subbasin for the purpose of participant group discussions in this project only. This decision was made to reflect the hydrologic connectivity of the mainstem in these sub-watersheds to Upper Klamath Lake, which drains directly into this portion of the mainstem and has a significant influence on it. This 'reorganization' will ensure that discussion of restoration priorities in this segment of the mainstem occurs in the same group discussing restoration priorities for Upper Klamath Lake.




# IFRMP - Redband Trout Range

## Legend:

-  Klamath Subbasin Groups
-  Redband Trout - current distribution <sup>1</sup>
-  Redband Trout - additional historical distribution <sup>1</sup>
-  Keno diversion area

## Data Sources:

1 - ODFW Oregon Fish Habitat Distribution Data (ODFW Data Clearinghouse):  
<https://nrimp.dfw.state.or.us/DataClearinghouse/default.aspx?p=202&XMLname=1167.xml>










## About This Map:


This map was produced as part of the Klamath Integrated Fisheries Restoration and Monitoring Plan (IFRMP) and is being used as the basis for determining range overlap of focal fish species with potential project locations during the restoration prioritization phase of this project (which is only one of several prioritization criteria). Find out more about this project at <http://kbifrm.psmfc.org/>.

Note that the Keno Diversion area shown on this map indicates reassignment of these sub-watersheds from the Lost Sub-basin to the Upper Klamath Lake subbasin for the purpose of participant group discussions in this project only. This decision was made to reflect the hydrologic connectivity of the mainstem in these sub-watersheds to Upper Klamath Lake, which drains directly into this portion of the mainstem and has a significant influence on it. This 'reorganization' will ensure that discussion of restoration priorities in this segment of the mainstem occurs in the same group discussing restoration priorities for Upper Klamath Lake.

# IFRMP - Shortnose Sucker Range

## Legend:

-  Klamath Subbasin Groups
-  Keno diversion area
-  Shortnose Sucker - current distribution <sup>1</sup>
-  Shortnose Sucker - additional historical distribution <sup>2</sup>
-  Shortnose Sucker - Designated Critical Habitat <sup>3</sup>
-  Shortnose Sucker - Designated Streams
-  Shortnose Sucker - Designated Lakes



## Data Sources:

- 1 - USFWS 2019 Species Status Assessment for the Endangered Lost River and Shortnose Sucker  
[https://www.researchgate.net/publication/333787292\\_Species\\_Status\\_Assessment\\_for\\_the\\_Endangered\\_Lost\\_River\\_Sucker\\_and\\_Shortnose\\_Sucker](https://www.researchgate.net/publication/333787292_Species_Status_Assessment_for_the_Endangered_Lost_River_Sucker_and_Shortnose_Sucker)
- 2 - Josh Rasmussen, USFWS, Personal communication.
- 3 - USFWS Critical Habitat Designations:  
<http://criticalhabitat.fws.gov/crithab>





## About This Map:

This map was produced as part of the Klamath Integrated Fisheries Restoration and Monitoring Plan (IFRMP) and is being used as the basis for determining range overlap of focal fish species with potential project locations during the restoration prioritization phase of this project (which is only one of several prioritization criteria). Find out more about this project at <http://kbifrm.psmfc.org/>.

Note that the Keno Diversion area shown on this map indicates reassignment of these sub-watersheds from the Lost Sub-basin to the Upper Klamath Lake subbasin for the purpose of participant group discussions in this project only. This decision was made to reflect the hydrologic connectivity of the mainstem in these sub-watersheds to Upper Klamath Lake, which drains directly into this portion of the mainstem and has a significant influence on it. This 'reorganization' will ensure that discussion of restoration priorities in this segment of the mainstem occurs in the same group discussing restoration priorities for Upper Klamath Lake.

# IFRMP - Summer Steelhead Range


## Legend:

-  Klamath Subbasin Groups
-  Summer Steelhead - current distribution<sup>1</sup>
-  Summer Steelhead - additional historical distribution<sup>1,2</sup>
-  Keno diversion area

## Data Sources:

1 - UC Davis - PISCES Fish Range and Occurrence Database: [https://pisc.es.ucdavis.edu/fish?term\\_node\\_tid\\_depth=All&term\\_node\\_tid\\_depth\\_1=All&field\\_species\\_commonname\\_value=](https://pisc.es.ucdavis.edu/fish?term_node_tid_depth=All&term_node_tid_depth_1=All&field_species_commonname_value=)

2 - ODFW Oregon Fish Habitat Distribution Data (ODFW Data Clearinghouse): <https://nrimp.dfw.state.or.us/DataClearinghouse/default.aspx?p=202&XMLname=1167.xml>



## About This Map:

This map was produced as part of the Klamath Integrated Fisheries Restoration and Monitoring Plan (IFRMP) and is being used as the basis for determining range overlap of focal fish species with potential project locations during the restoration prioritization phase of this project (which is only one of several prioritization criteria). Find out more about this project at <http://kbifrm.psmfc.org/>.





Note that the Keno Diversion area shown on this map indicates reassignment of these sub-watersheds from the Lost Sub-basin to the Upper Klamath Lake subbasin for the purpose of participant group discussions in this project only. This decision was made to reflect the hydrologic connectivity of the mainstem in these sub-watersheds to Upper Klamath Lake, which drains directly into this portion of the mainstem and has a significant influence on it. This 'reorganization' will ensure that discussion of restoration priorities in this segment of the mainstem occurs in the same group discussing restoration priorities for Upper Klamath Lake.

# IFRMP - Winter Steelhead Range

Steelhead?




## Legend:

-  Klamath Subbasin Groups
-  Winter Steelhead - current distribution<sup>1</sup>
-  Winter Steelhead - additional historical distribution<sup>1,2</sup>
-  Keno diversion area

## Data Sources:

1 - UC Davis - PISCES Fish Range and Occurrence Database: [https://pisces.ucdavis.edu/fish?term\\_node\\_tid\\_depth=All&term\\_node\\_tid\\_depth\\_1=All&field\\_species\\_commonname\\_value=](https://pisces.ucdavis.edu/fish?term_node_tid_depth=All&term_node_tid_depth_1=All&field_species_commonname_value=)

2 - ODFW Oregon Fish Habitat Distribution Data (ODFW Data Clearinghouse): <https://nrimp.dfw.state.or.us/DataClearinghouse/default.aspx?p=202&XMLname=1167.xml>



## About This Map:

This map was produced as part of the Klamath Integrated Fisheries Restoration and Monitoring Plan (IFRMP) and is being used as the basis for determining range overlap of focal fish species with potential project locations during the restoration prioritization phase of this project (which is only one of several prioritization criteria). Find out more about this project at <http://kbifrm.psmfc.org/>.

Note that the Keno Diversion area shown on this map indicates reassignment of these sub-watersheds from the Lost Sub-basin to the Upper Klamath Lake subbasin for the purpose of participant group discussions in this project only. This decision was made to reflect the hydrologic connectivity of the mainstem in these sub-watersheds to Upper Klamath Lake, which drains directly into this portion of the mainstem and has a significant influence on it. This 'reorganization' will ensure that discussion of restoration priorities in this segment of the mainstem occurs in the same group discussing restoration priorities for Upper Klamath Lake.